	Technical data sheet	Ref : FT.L0014an Page : 1/3
		Version date : 18/07/22

Glutamine – Penicillin - Streptomycin 100X

CAT N°: L0014

Storage conditions: -20°C

Shelf life: 24 months

Composition: Displayed on website, also available on request.

Colour: Colourless

pH: 4.7 ± 0.5

Osmolality: Not applicable

Endotoxin: <10 EU/ml

Sterility tests:

- Bacteria in aerobic and anaerobic conditions
- Fungi and yeasts

Cell Growth test: Not applicable


Other tests: Not applicable

Recommended use:

Use in cell culture applications at 10 ml/l. This concentration is for tissue culture media containing serum; serum-free media generally require lower concentration.

- Respect storage conditions of the product
- Do not use the product after its expiry date
- Store product in an area protected from light
- Manipulate the product in aseptic conditions (e.g.: under laminar air flow)
- Wear clothes adapted to the manipulation of the product to avoid contamination (e.g.: gloves, mask, hygiene cap, overall...)
- In order to preserve all product qualities, it is recommended to thaw out the flask, to aliquote, then to re-freeze the produced flasks rather than to thaw out and re-freeze the flask at each use.
- It is recommended to use the product immediately after its thaw out.

The product is intended to be used in vitro for research or further manufacturing only and not for use as an Active Pharmaceutical Ingredient or food or animal feed.

	Technical data sheet	Ref : FT.L0014an Page : 2/3
		Version date : 18/07/22

Application:

Antibiotics, combined with good sterile technique, help prevent microbiological contamination

When an irreplaceable culture becomes contaminated, determine if the contamination is bacteria, fungus, mycoplasma, or yeast. Isolate the contaminated culture from other cell lines. Clean incubators and laminar flow hoods with a laboratory disinfectant, and check HEPA filters.

Glutamine–Penicillin–Streptomycin solution at high concentration can be toxic to some cell lines; therefore, perform a dose response test to determine the level at which the solution becomes toxic.

The following is a suggested procedure for determining toxicity levels and decontaminating cultures.

- 1) Dissociate, count, and dilute the cells in antibiotic free medium. Dilute the cells to the concentration used for regular cell passage.
- 2) Dispense the cell suspension into a multiwell culture plate or several small flasks. Add the Glutamine–Penicillin–Streptomycin solution to each well in a range of concentrations.
- 3) Observe the cells daily for signs of toxicity such as sloughing, appearance of vacuoles, decrease in confluency, and rounding.
- 4) When the toxic level has been determined, culture the cells for two to three passages using the Glutamine–Penicillin–Streptomycin solution at a concentration one to two fold lower than the toxic concentration.
- 5) Culture the cells for one passage in an antibiotic-free media
- 6) Repeat step 4.
- 7) Culture the cells in antibiotic-free medium for 4 to 6 passages to determine if the contamination has been eliminated.

Uses:

Mode of Action: Penicillin acts by inhibiting bacterial cell wall synthesis.

Streptomycin inhibits prokaryote protein synthesis by preventing the transition from initiation complex to chain-elongating ribosome and causes miscoding.

Antimicrobial spectrum: Gram-negative and Gram-positive bacteria.

Potency of the penicillin in solution: 10 000 U/ml

Stability: 3 days at 37°C

Signs of Deterioration: Not applicable

Remarks: Not applicable



Technical data sheet

Ref : FT.L0014an
Page : 3/3

Version date : 18/07/22