



## G-418 (Geneticin) Solution

**CAT N°:** L0015

**Storage conditions:** - 20°C

**Shelf life:** 24 months

**Composition:** 50 mg/ml G-418 in water

**Colour:** Clear to very slightly hazy liquid

**pH:**  $5,2 \pm 0,5$

**Osmolality:**  $100 \pm 50$  mOsm/kg

**Endotoxin:** Non applicable

**Sterility tests:**

- bacteria in aerobic and anaerobic conditions
- fungi and yeast

**Cell Growth test:** Not applicable

**Other tests:** Not applicable

**Recommended use:** Dilute with buffer or cell culture medium

- Respect storage conditions of the product
- Do not use the product after its expiry date
- Store product in an area protected from light
- Manipulate the product in aseptic conditions (e.g.: under laminar air flow)
- Wear clothes adapted to the manipulation of the product to avoid contamination (e.g.: gloves, mask, hygiene cap, overall...)
- In order to preserve all product qualities, it is recommended to thaw out the flask, to aliquote, then to re-freeze the produced flasks rather than to thaw out and re-freeze the flask at each use.
- It is recommended to use the product immediately after its thaw out.

The product is intended to be used in vitro for research or further manufacturing only and not for use as an Active Pharmaceutical Ingredient or food or animal feed.

**Application:**

Antibiotic G418 is an aminoglycoside antibiotic similar in structure to gentamycin. It exhibits toxicity towards both eukaryotic and prokaryotic cells. The optimal concentration for selection and maintenance must be determined for each cell line. For bacteria and algae, concentrations of 5µg/ml or less are recommended. Animal cells may require up to 300-500 µg/ml. Typically, resistance is conferred by one of two dominant genes of bacterial origin, which can be expressed in eukaryotic cells. Cells that are multiplying will be affected sooner than those that are not.



## Technical data sheet

Ref : FT.L0015an

Page : 2/2

Version date : 28/11/2023

Cells in log phase may require three to seven days for selection. In general, concentrations of approximately 400 µg/ml for selection and 200 µg/ml for maintenance are required for mammalian cells.

**Uses:** Not applicable

**Signs of Deterioration:** Not applicable

**Remarks:** Not applicable