



Technical data sheet
Media Liquid

Ref : FT.L0640an
Page : 1/2

Version date : 02/03/26

NaCl salt solution 0.85%

CAT N°: L0640

Storage conditions: Room temperature, protected from light

Shelf life: 48 months

Composition: Displayed on website; also available on request

Colour: Clear solution

pH: For information

Osmolality: For information

Endotoxin: < 1 EU/ml

Sterility tests:

- Bacteria in aerobic and anaerobic conditions
- Fungi and yeasts

Cell Growth test: Not applicable

Other tests: Not applicable

Recommended use:

- Respect storage conditions of the product
- Do not use the product after its expiry date
- Store product in an area protected from light (not necessary for saline solutions).
- Manipulate the product in aseptic conditions (e.g.: under laminar air flow)
- Wear clothes adapted to the manipulation of the product to avoid contamination (e.g.: gloves, mask, hygiene cap, overall...)

The product is intended to be used in vitro for research or further manufacturing only and not for use as an Active Pharmaceutical Ingredient or food or animal feed.

Application:

The product is a saline solution at 0.85% prepared with water used for cell culture. The sodium chloride is a source of chlorine and sodium necessary to the production of components like sodium carbonate, sodium hydroxide or chlorate ions.

Uses:

The sodium chloride is widely used for biochemistry researches and molecular biology. This solution can be used for the realisation of phosphate buffer or SSC buffer (sodium chloride – sodium citrate), and also for the preparation of plasmidic DNA (removing of small fragments of nucleic acids) and for the precipitation of the DNA of samples containing SDS. The sodium



Technical data sheet
Media Liquid

Ref : FT.L0640an

Page : 2/2

Version date : 02/03/26

chloride is also used in the chemical sequencing of DNA. [1] and for protein crystallisation [2,3,4,5]

Signs of Deterioration:

Solution should be clear and free of particulate and flocculent material.

Do not use if solution is cloudy or contains precipitate.

Remarks:

[1] : Molecular Cloning: A Laboratory Manual, 3rd ed., Sambrook, J. F., et al., Cold Spring Harbor Laboratory Press (Cold Spring Harbor, NY: 2001), pp. 1.78-1.79, 2.71-2.75, 6.19-6.20, 6.24-6.25, 6.26, 12.23, 12.61-12.65, 16.43, A8.12.

[2] : Curtis, R. A., et al., Hydrophobic forces between protein molecules in aqueous solutions of concentrated electrolyte. *Biophys. Chem.*, **98(3)**, 249-265 (2002).

[3]: Smith, B. S., et al., Crystallization and preliminary X-ray analysis of ferric enterobactin receptor FepA, an integral membrane protein from *Escherichia coli*. *Acta Crystallogr. D Biol. Crystallogr.*, **54 (Pt 4)**, 697-699 (1998).

[4]: Discipio, R. G., et al., Crystallization of human complement component C5. *Acta Crystallogr. D Biol. Crystallogr.*, **54 (Pt 4)**, 643-646 (1998).

[5]: Solovyova, A., et al., Non-ideality by sedimentation velocity of halophilic malate dehydrogenase in complex solvents. *Biophys. J.*, **81(4)**, 1868-1880 (2001).